



Complete Summary

GUIDELINE TITLE

Micronutrient intake in children.

BIBLIOGRAPHIC SOURCE(S)

Caring for Australasians with Renal Impairment. Micronutrient intake in children. Nephrology 2005;10(Suppl 5):S213-14.

Hodson E. Micronutrient intake in children. Westmead NSW (Australia): CARI - Caring for Australasians with Renal Impairment; 2005 Dec. 4 p. [5 references]

GUIDELINE STATUS

This is the current release of the guideline.

COMPLETE SUMMARY CONTENT

SCOPE
METHODOLOGY - including Rating Scheme and Cost Analysis
RECOMMENDATIONS
EVIDENCE SUPPORTING THE RECOMMENDATIONS
BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS
IMPLEMENTATION OF THE GUIDELINE
INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT
CATEGORIES
IDENTIFYING INFORMATION AND AVAILABILITY
DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

- Chronic kidney disease (CKD)
- End-stage kidney disease (ESKD)

GUIDELINE CATEGORY

Assessment of Therapeutic Effectiveness
Management

CLINICAL SPECIALTY

Family Practice
Nephrology

Nutrition
Pediatrics

INTENDED USERS

Dietitians
Physicians

GUIDELINE OBJECTIVE(S)

To review the available evidence for the benefits and adverse effects of supplementation of vitamins and micronutrients in children with chronic kidney disease and end-stage kidney disease

TARGET POPULATION

Children with chronic kidney disease or end-stage kidney disease on dialysis

INTERVENTIONS AND PRACTICES CONSIDERED

1. Supplementation of water soluble vitamins and micronutrients
2. Supplementation of fat soluble vitamins (considered but not recommended)

MAJOR OUTCOMES CONSIDERED

- Rate of children meeting criteria for recommended daily intake of vitamins and micronutrients
- Dietary intake of micronutrients
- Serum levels of micronutrients

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

Databases searched: Medline (1996 to November Week 2 2003) and Embase (1980 to November 2003). Medical Subject Heading (MeSH) terms for kidney disease were combined with MeSH terms and text words for micronutrients. The Cochrane Renal Group Specialised Register of randomised controlled trials was also searched for relevant trials not indexed in Medline.

Date of searches: 1 December 2003.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Weighting According to a Rating Scheme (Scheme Given)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Levels of Evidence

Level I: Evidence obtained from a systematic review of all relevant randomized controlled trials (RCTs)

Level II: Evidence obtained from at least one properly designed RCT

Level III: Evidence obtained from well-designed pseudo-randomized controlled trials (alternate allocation or some other method); comparative studies with concurrent controls and allocation not randomized, cohort studies, case-control studies, interrupted time series with a control group; comparative studies with historical control, two or more single arm studies, interrupted time series without a parallel control group

Level IV: Evidence obtained from case series, either post-test or pretest/post-test

METHODS USED TO ANALYZE THE EVIDENCE

Review of Published Meta-Analyses
Systematic Review with Evidence Tables

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Expert Consensus

DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

Not stated

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

Comparison with Guidelines from Other Groups
Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Recommendations of Others. Recommendations regarding micronutrient intake in children with chronic kidney disease or end-stage renal disease on dialysis from the following groups were discussed: Kidney Disease Outcomes Quality Initiative, British Renal Association, Canadian Society of Nephrology, and European Best Practice Guidelines.

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

Definitions for the levels of evidence (I–IV) can be found at the end of the "Major Recommendations" field.

Guidelines

No recommendations possible based on Level I or II evidence

Suggestions for Clinical Care

(Suggestions are based on Level III and IV evidence)

- Supplements of water soluble vitamins are indicated in dialysis patients who are not receiving nutritional supplements.
- Supplements of vitamins A, B₁₂, and E are not indicated, since the dietary intake of these vitamins meets recommended daily intakes (RDIs) in children on dialysis.

Definitions:

Levels of Evidence

Level I: Evidence obtained from a systematic review of all relevant randomized controlled trials (RCTs)

Level II: Evidence obtained from at least one properly designed RCT

Level III: Evidence obtained from well-designed pseudo-randomized controlled trials (alternate allocation or some other method); comparative studies with concurrent controls and allocation not randomized, cohort studies, case-control studies, interrupted time series with a control group; comparative studies with historical control, two or more single arm studies, interrupted time series without a parallel control group

Level IV: Evidence obtained from case series, either post-test or pretest/post-test

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of supporting evidence is identified and graded for each recommendation (see "Major Recommendations").

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

Appropriate micronutrient intake in children with chronic kidney disease and end-stage kidney disease on dialysis

POTENTIAL HARMS

Not stated

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

Implementation and Audit

The number of children on dialysis who receive a vitamin supplement with or without nutritional supplements and the supplement used could be assessed through the paediatric renal dietitians and the Australian & New Zealand Paediatric Nephrology Association (ANZPNA).

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Living with Illness

IOM DOMAIN

Effectiveness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Caring for Australasians with Renal Impairment. Micronutrient intake in children. Nephrology 2005;10(Suppl 5):S213-14.

Hodson E. Micronutrient intake in children. Westmead NSW (Australia): CARI - Caring for Australasians with Renal Impairment; 2005 Dec. 4 p. [5 references]

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

2005 Dec

GUIDELINE DEVELOPER(S)

Caring for Australasians with Renal Impairment - Disease Specific Society

SOURCE(S) OF FUNDING

Industry-sponsored funding administered through Kidney Health Australia

GUIDELINE COMMITTEE

Not stated

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Author: Elisabeth Hodson

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

All guideline writers are required to fill out a declaration of conflict of interest.

GUIDELINE STATUS

This is the current release of the guideline.

GUIDELINE AVAILABILITY

Electronic copies: Available in Portable Document Format (PDF) from the [Caring for Australasians with Renal Impairment Web site](#).

Print copies: Available from Caring for Australasians with Renal Impairment, Locked Bag 4001, Centre for Kidney Research, Westmead NSW, Australia 2145

AVAILABILITY OF COMPANION DOCUMENTS

The following is available:

- The CARI guidelines. A guide for writers. Caring for Australasians with Renal Impairment. 2006 May. 6 p.

Electronic copies: Available from the [Caring for Australasians with Renal Impairment \(CARI\) Web site](#).

PATIENT RESOURCES

None available

NGC STATUS

This NGC summary was completed by ECRI Institute on March 31, 2008. The information was verified by the guideline developer on June 11, 2008.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 9/29/2008

